# **YEMEN**



## STATISTICAL PROFILE ON FEMALE GENITAL MUTILATION

Female genital mutilation (FGM) refers to "all procedures involving partial or total removal of the female external genitalia or other injury to the female genital organs for non-medical reasons." 1 While the exact number of girls and women worldwide who have undergone FGM remains unknown, at least 200 million girls and women have been cut in 30 countries with representative data on prevalence. FGM is a violation of girls' and women's human rights and is condemned by many international treaties and conventions, as well as by national legislation in many countries. Yet, where it is practised, FGM is performed in line with tradition and social norms to ensure that girls are socially accepted and marriageable, and to uphold their status and honour and that of the entire family. UNICEF works with government, civil society partners and other UN agencies towards the elimination of FGM in countries where it is still practised.

1. World Health Organization, Eliminating Female Genital Mutilation: An interagency statement, WHO, UNFPA, UNICEF, UNIFEM, OHCHR, UNHCR, UNECA, UNESCO, UNDP, UNAIDS, WHO, Geneva, 2008, p. 4.

2001

National decree/legislation banning FGM passed

### **SELECTED STATISTICS ON WOMEN'S STATUS**

9%	of women 20-24 years were married or in union before age 15
<b>32</b> %	of women 20-24 years were married or in union before age 18
<b>17</b> %	of women 20-24 years have given birth by age 18
49%	of women 15-49 years think that a husband/partner is justified in hitting or beating his wife/partner under certain circumstances
81%	of women 15-49 years make use of at least one type of information media at least once a week (newspaper, magazine, television or radio)


Source: DHS 2013







### **HOW WIDESPREAD IS THE PRACTICE?**

#### Almost one in five girls and women in Yemen have undergone FGM









### WHEN AND HOW IS FGM PERFORMED?

#### In Yemen, nearly nine out of ten girls who experienced FGM did so within their first week of life









Notes: Only categories with 25 or more unweighted cases are presented. Due to rounding, some of the data presented may not add up to 100 per cent. 'Health personnel' includes doctors, nurses, midwives and other health workers; 'Traditional practitioner' includes traditional circumcisers, traditional birth attendants, and other types of traditional practitioners.

Source for all charts on this page: DHS 2013

## WHAT ARE THE PREVAILING ATTITUDES TOWARDS FGM?


#### Around three in four girls and women think the practice should stop






## IS THE PRACTICE OF FGM CHANGING?


#### There has been practically no change in the prevalence of FGM







# **YEMEN**

## **INTER-COUNTRY STATISTICAL OVERVIEW**







Notes: Data on attitudes for Yemen refer to ever-married girls and women. In Liberia, girls and women who have heard of the Sande society were asked whether they were members; this provides indirect information on FGM since it is performed during initiation into the society. MICS data for Ghana (2011) could not be used to report on attitudes towards FGM due to the fact that information is missing for girls and women with no living daughters; data from MICS 2006 are used instead. In Liberia, only cut girls and women were asked about their attitudes towards FGM; since girls and women from practicing communities are more likely to support the practice, the level of support in this country as captured by DHS 2013 is higher than would be expected had all girls and women been asked their opinion. Prevalence data on FGM for girls and women aged 15 to 49 years and data on attitudes towards FGM are not available for Indonesia.

Updated January 2019

and RISKESDAS, 2004-2018.

Sources: DHS, MICS, Health Issues Survey, Population and Health Survey