

EGYPT



© UNICEF/NYHQ2005-1563/Pirozzi

STATISTICAL PROFILE ON FEMALE GENITAL MUTILATION

Female genital mutilation (FGM) refers to “all procedures involving partial or total removal of the female external genitalia or other injury to the female genital organs for non-medical reasons.”¹ While the exact number of girls and women worldwide who have undergone FGM remains unknown, at least 200 million girls and women have been cut in 30 countries with representative data on prevalence. FGM is a violation of girls’ and women’s human rights and is condemned by many international treaties and conventions, as well as by national legislation in many countries. Yet, where it is practised FGM is performed in line with tradition and social norms to ensure that girls are socially accepted and marriageable, and to uphold their status and honour and that of the entire family. UNICEF works with government and civil society partners towards the elimination of FGM in countries where it is still practised.

1. World Health Organization, *Eliminating Female Genital Mutilation: An interagency statement*, WHO, UNFPA, UNICEF, UNIFEM, OHCHR, UNHCR, UNECA, UNESCO, UNDP, UNAIDS, WHO, Geneva, 2008, p. 4.

2008

National legislation criminalizing FGM passed

SELECTED STATISTICS ON WOMEN’S STATUS

2% of women 20-24 years were married or in union before age 15

17% of women 20-24 years were married or in union before age 18

7% of women 20-24 years have given birth by age 18

36% of ever-married women 15-49 years think that a husband/partner is justified in hitting or beating his wife/partner under certain circumstances


97% of ever-married women 15-49 years make use of at least one type of information media at least once a week (newspaper, magazine, television or radio)

Source: DHS 2014


HOW WIDESPREAD IS THE PRACTICE?

FGM is nearly universal among girls and women of reproductive age in Egypt


Percentage of girls and women aged 15 to 49 years who have undergone FGM, by region




Percentage of girls and women aged 15 to 49 years who have undergone FGM, by residence and wealth quintile



Percentage of girls aged 1 to 14 years who have undergone FGM (as reported by their mothers), by residence, mother's education and household wealth quintile




Among daughters of cut girls and women, percentage of girls aged 1 to 14 years who have undergone FGM (as reported by their mothers), by mothers' attitudes about whether the practice should continue



WHEN AND HOW IS FGM PERFORMED?

Three-quarters of girls who underwent FGM were cut by health personnel

Percentage distribution of girls aged 15 to 19 years who have undergone FGM, by age at which cutting occurred



Percentage distribution of girls aged 1 to 14 years who have undergone FGM/C (as reported by their mothers), according to the type of person/practitioner performing the procedure




Notes: The boundaries and the names shown and the designations used on the map do not imply official endorsement or acceptance by the United Nations. Only categories with 25 or more unweighted cases are presented. Due to rounding, some of the data presented may not add up to 100 per cent. Prevalence data of Frontier Governorates do not include North and South Sinai. Prevalence data for girls aged 1 to 14 reflect their current, but not final, FGM status since some girls who have not been cut may still be at risk of experiencing the practice once they reach the customary age for cutting. Therefore, the data on prevalence for girls under age 15 is actually an underestimation of the true extent of the practice. Since age at cutting varies among settings, the amount of underestimation also varies and this should be kept in mind when interpreting all FGM prevalence data for this age group. 'Health personnel' includes doctors, nurses, midwives and other health workers; 'Traditional practitioner' includes traditional circumcisers, traditional birth attendants, traditional midwives, other types of traditional practitioners and dayas, ghagarias and barbers. Questions on type of cutting only differentiated infibulation from non-infibulating forms of FGM. Information on type of cutting was not asked in more recent surveys in Egypt.

Source for all charts on this page: Health Issues Survey 2015


WHAT ARE THE PREVAILING ATTITUDES TOWARDS FGM?

Around half of people in Egypt support the continuation of FGM and believe it is a religious requirement


Percentage distribution of girls and women aged 15 to 49 and boys and men aged 15 to 49 years who have heard about FGM, by their attitudes about whether the practice should continue



Percentage of girls and women aged 15 to 49 years and boys and men aged 15 to 49 years who have heard of FGM/C and believe the practice is required by religion



Percentage of girls and women aged 15 to 49 years who have heard about FGM and think the practice should continue, by residence, education and age




Source for all of the above charts: Health Issues Survey 2015


IS THE PRACTICE OF FGM CHANGING?

Fewer adolescent girls have experienced FGM compared to older women, and support for FGM has declined

Percentage of girls and women aged 15 to 49 years who have undergone FGM, by current age



Percentage of ever-married girls and women aged 15 to 49 years who have undergone FGM, and percentage ever-married of girls and women aged 15 to 49 years who have heard about FGM and think the practice should continue



Source: Health Issues Survey 2015


Note: Data from the DHS 2008 and Health Issues Survey 2015 were recalculated for ever-married girls and women to allow comparison with earlier surveys which only collected data on FGM from ever-married girls and women.

EGYPT




INTER-COUNTRY STATISTICAL OVERVIEW


Percentage of girls and women aged 15 to 49 years who have undergone FGM



Percentage of girls aged 0 to 14 years who have undergone FGM (as reported by their mothers)



Percentage of girls and women aged 15 to 49 years who have heard about FGM and think the practice should continue



Notes: Data on attitudes for Yemen refer to ever-married girls and women. In Liberia, girls and women who have heard of the Sande society were asked whether they were members; this provides indirect information on FGM since it is performed during initiation into the society. Egypt data refer to girls aged 1 to 14 years and Indonesia data refer to girls aged 0 to 11 years. An older source is used to report on the prevalence of FGM among girls aged 0 to 14 years for Gambia (MICS 2010) and Uganda (DHS 2011) since the latest source did not collect these data. MICS data for Ghana (2011) could not be used to report on attitudes towards FGM due to the fact that information is missing for girls and women with no living daughters; data from MICS 2006 are used instead. In Liberia, only cut girls and women were asked about their attitudes towards FGM; since girls and women from practicing communities are more likely to support the practice, the level of support in this country as captured by DHS 2013 is higher than would be expected had all girls and women been asked their opinion. Prevalence data on FGM for girls and women aged 15 to 49 years and data on attitudes towards FGM are not available for Indonesia.